

Gabapentin Rosemont Oral Solution

50mg/mL

Instructions for administration via nasogastric (NG) or percutaneous endoscopic gastrostomy (PEG) tubes.

Gabapentin Rosemont Oral Solution is suitable for use with the following type of NG and PEG tubes:

Material	External Bore Size (Fr Unit)	Internal Diameter (mm)	Maximum Length (cm)	Recommended Flush Volume (mL)
Silicone	4	0.8	125	5mL
	6	1.0	125	5mL
	10	2.0	125	10mL
PVC	4	0.8	125	5mL
	8	1.5	125	5mL
	12	2.5	125	10mL
Polyurethane	4	0.8	125	5mL
	8	1.5	125	5mL
	12	2.6	125	10mL
	18	4.0	75	10mL

Ensure that the enteral feeding tube is free from obstruction before administration.

1. Flush the enteral tube with water, using the minimum flush volume required.

2. Administer the required dose of Gabapentin Rosemont Oral Solution with a suitable measuring device.

3. Flush the enteral tube with water again using the minimum flush volume required.

This product should be administered with silicone, PVC, polyurethane NG or PEG tubes **only**.

Healthcare professionals should be aware that with air flushing procedure there is a risk of under dosing (up to 50%). It is therefore recommended that only water flush is used.

The oral syringe included in the pack is only for patients who are able to swallow the medicine. HCPs must use another suitable device.

Any unused product or waste material should be disposed of in accordance with local requirements.



DTM060.1 July 2020

Rosemont Pharmaceuticals Ltd. Rosemont House, Yorkdale Industrial Park, Braithwaite Street, Leeds LS11 9XE T +44 (0)113 244 1400 F +44 (0)113 245 3567 E infodesk@rosemontpharma.com Sales/ Customer Service: T +44 (0) 113 244 1999 F +44 (0) 113 246 0738 W www.rosemontpharma.com

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard Adverse events should also be reported to Rosemont Pharmaceuticals Ltd on 0113 244 1400.